

GRB 081203A: Swift-UVOT captures the earliest ultraviolet spectrum of a Gamma Ray Burst

N. P. M. Kuin^{1*}, W. Landsman², M. J. Page¹, P. Schady¹, M. Still¹,
A. A. Breeveld¹, M. De Pasquale¹, P. W. A. Roming³, P. J. Brown³,
M. Carter¹, C. James¹, P. A. Curran¹, A. Cucchiara³, C. Gronwall³,
S. T. Holland², E. A. Hoversten³, S. Hunsberger³, T. Kennedy¹, S. Koch³,
H. Lamoureux¹, F. E. Marshall², S. R. Oates¹, A. Parsons², D. M. Palmer⁴,
and P. J. Smith¹

¹Mullard Space Science Laboratory/UCL, Holmbury St. Mary, Dorking, Surrey, RH5 6NT, UK

²NASA/Goddard Space Flight Center, Greenbelt, MD 20771, USA

³Department of Astronomy & Astrophysics, Penn State University, 525 Davey Laboratory, University Park, PA 16802, USA

⁴Los Alamos National Laboratory, P.O. Box 1663, Los Alamos, NM 87545

Accepted Received: Revision 1

ABSTRACT

We present the earliest ultraviolet spectrum of a gamma-ray burst (GRB) as observed with the *Swift*-UVOT. The GRB 081203A spectrum was observed for 50 seconds with the UV-grism starting 251 seconds after the *Swift*-BAT trigger. During this time the GRB was ≈ 13.4 mag (*u*-filter) and was still rising to its peak optical brightness. In the UV-grism spectrum we find a damped Ly- α line, Ly- β , and the Lyman continuum break at a redshift $z = 2.05 \pm 0.01$. A model fit to the Lyman absorption implies a gas column density of $\log N_{\text{HI}} = 22.0 \pm 0.1 \text{ cm}^{-2}$, which is typical of GRB host galaxies with damped Ly- α absorbers. This observation of GRB 081203A demonstrates that for brighter GRBs ($v \approx 14$ mag) with moderate redshift ($0.5 < z < 3.5$) the UVOT is able to provide redshifts, and probe for damped Ly- α absorbers within 4–6 minutes from the time of the *Swift*-BAT trigger.

Key words: gamma-rays: bursts - techniques: spectroscopy - instrumentation: spectrographs

1 INTRODUCTION

Gamma-ray bursts (GRBs) are the most luminous cosmic explosions known, and their afterglows extend throughout the electromagnetic spectrum, from X-ray to radio wavelengths. Spectra of GRB afterglows in the ultraviolet (UV) to optical wavelength range are particularly important for the study of GRBs, because the spectral positions of absorption lines from the surrounding environment provide the redshifts, and hence the distances and luminosities of GRBs. Under exceptionally favourable conditions, prompt, very near-UV GRB spectra have been obtained from the ground (i.e. *VLT*-UVES, which in principle is sensitive to wavelengths longward of 3000 Å, observed a GRB in ≈ 10 minutes). However, such observations are rare due to atmo-

spheric limitations, time of occurrence, and the source position in the sky. Shorter wavelength UV GRB spectra have so far only been taken with the *HST*/STIS (Smette et al. 2001), at least 3 days after the GRB trigger. In this letter we present the first prompt UV spectrum of a GRB. The spectrum was obtained with the UV-grism of the *Swift* Ultraviolet/Optical Telescope (UVOT; Roming et al. 2005).

Since the launch of the *Swift* satellite (Gehrels et al. 2004) in 2004, the UVOT has been making prompt photometric observations of GRBs in the optical and UV, simultaneously with the X-ray and gamma-ray observations taken by the X-Ray Telescope (XRT; Burrows et al. 2005) and the Burst Alert Telescope (BAT; Barthelmy et al. 2005), respectively. The UVOT photometric system (Poole et al. 2008) provides photometry in three UV and three optical bands, and in one *white*-filter which is sensitive over the range of 1700–8000 Å.

* email: npmk@mssl.ucl.ac.uk




Figure 1. The lightcurve of GRB 081203A shows that the spectrum was taken during the rise. The photometry point derived from the spectrum, indicated with an arrow, has a larger uncertainty than those of the lenticular filters.

Two grisms are also included in the UVOT filter wheel to enable spectra of GRBs to be obtained. Until recently they have not been used in the GRB observing sequence because early in the mission the data indicated that most GRB afterglows were too faint for grism observations, even at early times (Roming et al. 2006). However, after four years of *Swift* observations, our GRB sample is large enough that we can meaningfully reassess the frequency of GRB afterglows which are bright enough for grism spectroscopy, and ascertain the optimum timing for their inclusion in the automated observing sequence. The catalog of GRB afterglows observed with the UVOT in the first two-and-a-half years (Roming et al. 2008) shows that UVOT observes ≈ 90 each year, of which ≈ 25 are detected by UVOT in at least one exposure. In its first four years of operation, *Swift* observed 5 GRBs (GRB 050525A, GRB 050922C, GRB 061007, GRB 080319B and GRB 080810) brighter than 14th magnitude in *v*. It is GRBs like these that should provide good spectra through the UVOT grisms.

The UV-grism was introduced into the automated UVOT GRB response on the 7th Oct, 2008. In this latest sequence, the grism spectrum is 50 seconds long, preceded by an initial 150 second *white*-filter finding chart and followed by a second finding chart in the *u* filter. With a median *Swift* slew time of 86 seconds (Roming et al. 2008), the grism observation typically begins 225-275 seconds after the burst trigger, and covers the wavelength range from ≈ 1700 -5800 Å. The onboard software autonomously replaces the grism exposure with a *u*-filter exposure for bursts which are faint in the BAT, and therefore unlikely to be viable grism targets.

In the following sections we present the first grism image of a GRB taken with the UVOT UV-grism, the resulting spectrum of GRB 081203A, and its corresponding analysis.

2 OBSERVATIONS AND DATA REDUCTION

The BAT instrument triggered on GRB 081203A at $T = 2008$ -12-03 at 13:57:11 UT (Parsons et al. 2008). The mask-weighted light curve showed two overlapping peaks: the first starts at T -69 seconds and peaks at T +10 seconds, the second peaks at T +32 seconds and ends at T +405 seconds. UVOT started settled observations of GRB 081203A 93 seconds after the trigger beginning with a 150 second *white*-filter finding chart, followed by a 50 second UV-grism exposure starting 251 seconds after the trigger, a second finding chart in *u*, and then exposures in the other UVOT photometric filters (De Pasquale & Parsons 2008). The UV-grism spectrum was taken during the decay of the second BAT gamma-ray peak. The best source position was determined by the UVOT, which located a fading source at $RA_{J2000} = 15^h 32^m 07.58s$ ($=233.03158deg$), $Dec_{J2000} = +63^d 31^m 14.9s$ ($=63.52081deg$), with an uncertainty of $0.5''$ (90% confidence limit; De Pasquale & Parsons 2008), consistent with the refined XRT position (Goad et al. 2008). Using the grism spectrum, a provisional redshift of 2.1 was reported by Landsman et al. (2008). The X-ray spectrum, extracted from the XRT Window Timing data, overlaps with the grism exposure and has an exposure time of 558 s, starting 87 seconds after the trigger (La Parola et al. 2008).

The source was detected in the *u*, *b*, *v*, *white*, *uvw1* filters, and only faintly in the *uvw2* filter. It was not detected in the *uvm2* filter, which is consistent with the filter response curves and the shape of the UV-grism spectrum. The brightness of the GRB can be measured from the zeroth order of the grism image, which provides a magnitude of $b = 13.76 \pm 0.23$ at the time of the grism exposure. Folding the extracted spectrum (see Section 3) through the *u* filter response leads to a magnitude of 13.4 mag during the grism exposure with a systematic error estimated to be 25%. The light curve of GRB 081203A is shown in Figure 1.

A cutout of the grism image with the GRB spectrum is shown in Figure 2. The zeroth order of a 15th magnitude star, falls on the GRB spectrum around 5000 Å. The image was processed using the HEADAS-6.5.1 software to remove the MODULO-8 pattern (Poole et al. 2008).

The raw count rate spectrum was extracted using the UVOTIMGRISM program. The data were binned up by a factor of three, which reduces any correlations caused by the image rotation during the spectral extraction but keeps the bin size smaller than the wavelength resolution. The wavelength scale was adjusted to the latest UV-grism wavelength calibration (to be released shortly) for the nominal filter wheel position, and is accurate to 15 Å. The UV-grism spectrum has a resolution of $R \approx 150$ at 2600 Å, with a dispersion of ~ 3.2 Å/pixel. It can be seen that blueward of 2780 Å no signal above the background is seen. Although wavelengths in the first order spectrum longer than 2850 Å can be contaminated by a second order spectra, this is not a concern here since there is no source flux shortward of 2780 Å, thus, the second order spectrum will only affect the first order for wavelengths longer than 5549 Å.

3 SPECTRAL MODELLING

The background subtracted spectrum of GRB 081203A is provided in Figure 3, overlaid by our best-fit spectral




Figure 2. The UV-grism image of GRB 081203A. The GRB first order spectrum is inside the black rectangle. The zeroth order is indicated by a black circle. The ‘X’ in the top left corner indicates the contaminating zeroth order of a star.

model. Calibrations for the wavelength scale, line spread function, and telescope throughput have been extracted from a development version of the Swift calibration data base. Wavelengths are currently good to 0.5% systematic accuracy (1σ) and the flux calibration is good to an estimate of 25%.

Neglecting the systematic uncertainty in the wavelength scale, Figure 3 reveals a broad absorption line at $3706 \pm 11 \text{ \AA}$ that we identify with the $\text{Ly}\alpha$ transition of hydrogen, and a corresponding Lyman continuum edge at $2779 \pm 8 \text{ \AA}$, shortward of which no significant source detection is made. $\text{Ly}\beta$ is weakly detected at $3126 \pm 8 \text{ \AA}$. Wavelength errors and the best-fit parameters are at the 1σ level.

Within the spectral model, the afterglow continuum is characterised by a single power law attenuated by neutral gas and dust from both the Milky Way and host galaxy of the burst. Both galactic and host galaxy continuum absorption at wavelengths shorter than the Lyman edge are modelled using the photo-electric cross-sections of Morrison and McCammon (1983) and the relative element abundances of Anders & Ebihara (1982). The Milky Way extinction law has been taken from the analytical description of Pei (1992), and the host extinction is assumed identical to the Small Magellanic Cloud law, also from Pei (1992). Host galaxy hydrogen Lyman series absorption from $n = 1-100$ is modelled following the algorithm described in Totani et al. (2006) and based upon Peebles (1991). Oscillator strengths for transitions $n = 1-31$ are taken from Morton (1991) and extrapolated to $n \leq 100$ thereafter. The wavelength-dependent optical depth calculated within the model is also broadened by a macroscopic velocity field within the absorbing column. We assume a Gaussian velocity field, characterised by its full-width half-maximum (FWHM) velocity.

The combined model contains four free parameters: cosmological redshift (z), neutral hydrogen column density (N_{HI}), FWHM of the absorbers velocity field (v), and the normalization of the afterglow continuum. In order to constrain the continuum emission over the relatively narrow spectral range of the UV-grism, five additional model pa-




Figure 3. The UV-grism first order spectrum of GRB 081203A. The model shown here as a red line includes the Lyman forest as discussed in Section 3.

rameters are pre-determined. The Galactic column density towards the burst, $\log N_{\text{H,g}} = 20.23$, is taken from the map of Kalberla et al. (2005). Galactic dust is quantified by the extinction coefficient $A_{\text{V,g}} = 0.06$, taken from Schlegel et al. (1998). The continuum spectral index ($\beta = 0.90 \pm 0.01$), the neutral column density within the host galaxy ($\log N_{\text{H,h}} = 21.7 \pm 0.1$), and the host galaxy dust extinction coefficient ($A_{\text{V,h}} = 0.08$) were all measured by a fit to the broad band spectral energy distribution (SED) of the afterglow, interpolated to the $T+700\text{s}$ epoch, and incorporating six lenticular filter magnitudes from the UVOT and the X-ray spectrum from the XRT. This method is described by Schady et al. (2007). The detailed fit to the SED will be provided in a forthcoming paper concerning the spectral and temporal behaviour of GRB 081203A.

The model fit to the grism spectrum was performed using XSPEC 12.5.0 (Arnaud 1996). The best-fit provides $\chi^2 = 301.9$ for 231 degrees of freedom (dof), $\log N_{\text{HI}} = 22.09^{+0.09}_{-0.12}$, $z = 2.046 \pm 0.002$, and $v < 598 \text{ km/s}$ (90% confidence limit). The significance of the line detections was determined by freezing the redshift. The fit to the spectrum in the $3400 - 4000 \text{ \AA}$ wavelength range gives a 4.0σ detection of $\text{Ly}\alpha$, and $\text{Ly}\beta$ is detected at the 1.5σ level in the $3000 - 3300 \text{ \AA}$ wavelength range. The best-fit column density satisfies the definition for a damped $\text{Ly}\alpha$ absorption system (DLA). Taking into account systematic errors in the wavelength dispersion, $z = 2.05 \pm 0.01$.

At a redshift of 2.05, we expect some contamination in the spectrum from absorption systems located between the Earth and the host (Madau 1995). Within the limits of the signal-to-noise, there is no compelling evidence for discrete $\text{Ly}\alpha$ absorption from intervening galaxies. We do however investigate whether an undetected, or unresolved, Lyman forest could have an impact on the measured properties. We add the statistically-based algorithm characterising the cosmological hydrogen density from Madau (1995) to the model, which effectively absorbs photons blueward of the host galaxy $\text{Ly}\alpha$ line (Figure 3). The best fit for this situation provides $\chi^2 = 271.4$ for 231 dof, $\log N_{\text{HI}} = 22.02^{+0.11}_{-0.12}$, $z = 2.05 \pm 0.01$, and $v < 474 \text{ km/s}$. The Madau model gives a slightly better fit to the data.




Figure 4. The figure shows the number of GRBs with an associated damped Lyman- α system (DLA). The left panel shows that the N_{HI} -column density of GRB 081203A as determined in this paper is quite typical. The right-hand panel shows that the redshift of GRB 081203A is one of the lowest measured for a GRB with a DLA. The UVOT is capable of measuring redshifts below $z = 1.9$ that cannot be measured from the ground. The data for this figure were taken from the literature (Prochaska et al. 2007; Jakobsson et al. 2006; Savaglio 2008).

4 DISCUSSION AND CONCLUSIONS

With the rapid and precise locations of GRBs available with *Swift*, the number of GRBs available with good signal-to-noise spectroscopic data has doubled. Of these, 16% have an associated DLA system (Prochaska et al. 2007; Jakobsson et al. 2006). The column density of $N_{HI} = 1.04 \times 10^{22} \text{ cm}^{-2}$ in GRB 081203A is typical of GRB-DLAs (Prochaska et al. 2008), but its redshift is amongst the smallest of the sample (see Figure 4). It also has the lowest redshifts observed in a GRB for which both the Lyman break and Ly- α have been obtained.

Since it is possible for UVOT spectra to be taken during the rising phase of the afterglow, and while the gamma-ray emission is still in progress, as was the case for GRB 081203A, useful constraints on the evolution of the ionization state of the gas in the GRB environment may be obtained by comparison to ground-based spectra taken later in the decay phase of the afterglow. Getting an early spectrum with UVOT opens up the opportunity to increase the sample of *Swift* GRBs with redshifts. As well as its obvious advantage for UV coverage, the UVOT is capable of taking spectra of GRBs with positions that are unobtainable for ground based observations. Indeed, GRB 081203A was a twilight object for ground based observers, and we are not aware of any ground-based spectroscopic follow-up of this object.

We have shown that it is possible to obtain useful spectra using the UVOT UV-grism in the automated *Swift* response to new GRBs. The combined measurement of the Ly-continuum edge, Ly- α , and possibly Ly- β provides a good measurement of the redshift in the UVOT UV-grism spectra for redshifts of GRBs between $0.5 < z < 3.5$. Using such spectra, we expect to find new DLAs in the range

$0.5 < z < 2.0$. *Swift* observations can in principle probe the different components of the interstellar medium by providing a measurement of HI from the grism spectrum, dust extinction from the photometry, and metallicity from the X-rays.

ACKNOWLEDGEMENTS

This work was supported by the UK Science and Technology Facilities Council through a grant for *Swift* Post Launch Support at UCL-MSSL. This work is sponsored at PSU by NASA contract NAS5-00136. We acknowledge useful comments by the anonymous referee which led to improvement of the grism analysis. We would like to dedicate this letter to the late Richard Bingham, whose visionary optical design for the UVOT grism made these observations possible.

REFERENCES

- Anders, E. and Ebihara, M., 1982, *Geochimica et Cosmochimica Acta*, 46, 2363
- Arnaud, K.A., 1996, in Jacoby G. and Barnes J., eds., *ASP Conf. Ser. Vol. 101, Astronomical Data Analysis Software and Systems V*, p. 17
- Barthelmy, S. *et al.*, 2005, *Space Sci. Rev.* 120, 143
- Burrows, D. *et al.*, 2005, *Space Sci. Rev.* 120, 165
- De Pasquale, M. and Parsons, A., 2008, *GCN Circ. No.* 8603
- Gehrels, N., *et al.*, 2004, *ApJ*, 611, 1005
- Goad, M. *et al.*, 2008, *GCN Circ. No.* 8598
- Jakobsson, P. *et al.*, 2006, *A&A*, 460, L13
- Kalberla P. M. W., *et al.*, 2005, *A&A*, 440, 775
- Landsman, W.B. *et al.*, 2008, *GCN Circ. No.* 8601
- La Parola, V. *et al.*, 2008, *GCN Circ. No.* 8609
- Madau, P., 1995, *ApJ*, 441, 18
- Mason, K.O., *et al.*, 2004, *Proc. SPIE* 5165, 277
- Morrison, R. and McCammon, D., 1983, *ApJ* 270, 119
- Morton D.C., 1991, *ApJS*, 77, 119
- Parsons, A. *et al.*, 2008, *GCN Circ. No.* 8595
- Peebles P.J.E. 1993, *Principles of Physical Cosmology*, Princeton University Press
- Pei, Y.C., 1992, *ApJ*, 395, 130
- Poole, T., *et al.*, 2008, *MNRAS* 383, 627
- Prochaska, J.X. *et al.*, 2007, *ApJ*, 666, 267
- Prochaska, J.X. *et al.* 2008, *ApJ*, 685, 344
- Roming, P.W.A. *et al.*, 2005, *Space Sci.Rev.* 120, 95
- Roming, P.W.A. *et al.*, 2006, *ApJ*, 652, 1416
- Roming, P.W.A. *et al.*, 2009, *ApJ*, 690, 163
- Savaglio, S., 2008, private communication
- Schady, P. *et al.*, 2007, *MNRAS* 377, 273
- Schlegel, D.J., Finkbeiner, D.P., Davis, M., 1998, *ApJ* 500, 525
- Smette, A. *et al.*, 2001, *ApJ*, 556, 70
- Totani, T. *et al.* 2006, *PASJ* 58, 485
- Ukwatta, T.N., *et al.*, 2008, *GCN Circ. No.* 8599